

Working with young people who are bullied: Tips for mental health professionals

Mental health professionals are often asked to support young people who have been bullied. It is important to know what these young people need from us- and what approaches are unlikely to help.

Children Who Are Bullied Need:

- permission and support to tell what has happened to them and to talk about their feelings.
- protection from continued bullying through adult supervision, consequences for the young people who bully, and adults' taking reports of bullying seriously;
- strong, positive relationships with adults and peers;
- assistance from peers in feeling that they belong;
- assistance in not blaming themselves for the bullying.
- They also may need support with post-traumatic stress symptoms even after the bullying has been stopped.

What Probably Won't Help:

- The primary focus of our interventions should be: (1) protection from repeat bullying, and (2) limiting the effects of past bullying. If mental health professionals focus primarily on how these young people should prevent future bullying through their own actions, we risk confirming their belief that the bullying was their fault.
- Encouraging children who have been bullied to 'work things out' with the young person who has bullied them (especially alone or with the support of peer mediators) may well make things worse. Mediation-based approaches tend to imply that both parties are partly to blame. Young people who bully are unlikely to listen to the youth they have chosen as their targets. When mental health professionals encourage such discussions, we risk that the young person will be further victimized or controlled.
- Avoid advice that will not work or that they have already found ineffective. When we ask young people to 'tell them how you feel;' 'just walk away;' 'just not let it bother them' or 'just make a joke of it,' we may risk them feeling that they have failed when these strategies don't work. If they have already tried these strategies unsuccessfully before talking with us, we diminish their regard for us as resources.
- Be careful not to treat these youth like time bombs at imminent risk of aggression to others, unless there are indications of such risk. Most young people who are bullied direct their anger at themselves, rather than at others.

Why young people who have been bullied may not talk about it:

Young people who have been bullied may resist talking about the bullying, for several reasons:

- They may be ashamed of themselves. The person who bullied them may have succeeded in convincing them that they deserve the bullying.
- They may have come to believe the usually misguided message that if they only 'stood up to' the bully once, the bullying would stop. As a result, they may feel weak because they were not able to stop the bullying on their own.
- They may have found that telling adults does no good- that they are blamed, not believed, told to stop tattling, or given advice that doesn't work.

Counseling young people who are bullied

- They may not believe that adults will protect them. They may have found that they will not be protected from retaliation for telling.
- They may have become convinced that telling an adult about a problem is a sign of immaturity or that it is not “macho” (for boys).
- They may not be convinced that adults care about stopping bullying, because they may have seen adults watching bullying without doing anything..

Helping these youth talk about the bullying and avoid self-blame:

To encourage these youth to talk about what has happened requires the same orientation we would use with targets of sexual harassment, spouse abuse, or racism- three behaviors that share the same dynamics as bullying:

- The victim did not cause the problem and will not be asked to solve the problem alone;
- The victim has the right to tell, to be protected from retaliation for telling, and to have advocacy and protection. The person causing the abuse will be held fully responsible for his/her behavior.
- The victim may have been hurt in stages- by the abuse, by the inaction of others who should have helped, by unearned self-blame, and by social isolation. Each of these issues should be addressed.

How can mental health professionals effectively intervene to help a young person who has been bullied?

- Advocate- and help parents advocate for- supervision, staff training, consequences for the youth who bullies, and adult and peer support in the situation where bullying is taking place.
- Work with parents, school, and (through the school, coach, or other situation) with peers, to help the young person be less isolated and more connected with peers.
- Help parents and youth lessen isolation through connecting in friendship with peers who have shared interests.
- Explore unearned self-blame and help the young person see that the child who bullied him or her is responsible for the bullying.
- Explore and commend the steps the young person used to try to stop the bullying. Help him or her find effective ways to be safe. Identify advocates and resources in the environment. Help the student accept the probable benefits of using these allies.
- Assist the youth in reducing the pain associated with memories of victimization. The techniques that mental health professionals use to help a young person experiencing loss and anxiety will be helpful.
- Remember that young people who are bullied may also bully others, though most do not. Mental health professionals may also need to focus on aggression toward others.

What about young people who seem to provoke bullying?

For the most part, these young people have few other ways to make connections with peers. They may find that provoking peers' aggression toward them is better than being ignored. As we help them make more positive peer connections they can often give up the provoking behaviors. This process may take some time. During this transition the young people who seem to be provoking the bullying, and anyone who bullies them, should *both* be held fully responsible for their actions.

Counseling young people who are bullied

Often both youth involved will earn consequences in such a situation. It will not work to overlook harassment by peers because we think it is justified by the provoking behavior.

What about young people who over-react to bullying?

Some young people cry, yell, or otherwise react strongly to small acts of bullying, strongly reinforcing the person who bullies. [Might it be possible to provide an example of over-reaction?] These young people need to learn to moderate their reactions to bullying. Mental health professionals can teach them to calm themselves and to focus on the individual incident rather than to be overwhelmed by what 'always' happens. At the same time these young people often need to have specific adult advocates in their environment who will make sure that they do not get too much sympathy **and** that their reports of bullying will be taken seriously. Sometimes parents are part of the problem if they over-react themselves. These parents are often able to moderate their own overprotective reactions when they know these advocates will make sure the child's concerns are heard.

What about relational aggression?

If the bullying has taken place over time within a relationship that the young person describes as a friendship, mental health professionals should work with the young person and his or her parents to help them disconnect from that relationship in the same way that we would help someone disconnect from an abusive dating relationship. Children and youth should not be encouraged to 'work things out' with the abusive 'friends.'

In summary.....

Mental health professionals can help stop bullying as they advocate for effective interventions and help parents be more effective advocates. They can help young people overcome the emotional aftermath of bullying after supports and protection are in place.

References and resources:

- Davis, S. (2005). *Schools Where Everyone Belongs: Practical Strategies for Reducing Bullying* Champaign, IL, USA, Research Press
- Juvonen, J., Graham, S., Schuster, M. (2003). Bullying among young adolescents: The strong, the weak, and the troubled. *Pediatrics*, 112: 1231-1237
- McCoy, E. (1997). *What to Do... When Kids Are Mean to Your Child*. Pleasantville, NY: Reader's Digest.
- Mullin-Rindler, Nancy (2003). Relational aggression and bullying: It's more than just a girl thing. Center for Research on Women, Wellesley MA. *Wellesley Centers for Women Working Paper Series*, 2003. Working Paper # 408.
- Mynard, H., Joseph, S., & Alexander, J. (2000). Peer victimization and post-traumatic stress in adolescence. *Personality and Individual Differences*, 29, 815-821.
- National Association of School Psychologists. (2002) Bullying prevention: what schools and parents can do. *National Association of School Psychologists*, Bethesda, MD.
http://www.naspcenter.org/resourcekit/bullying_new_rk.html

Counseling young people who are bullied

New South Wales Health Department (2001) "What parents should know about bullying"
<http://www.mhcs.health.nsw.gov.au/health-public-affairs/mhcs/publications/5705.html>

Olweus, D. (1993). *Bullying at school: What we know and what we can do*. Oxford, UK: Blackwell

Rigby, K. (2000). Effects of peer victimisation in schools and perceived social support on adolescent well-being. *Journal of Adolescence*, 23, 57-68.

Sanford, L. (1992). *Strong At The Broken Places*. New York: Avon.

Starr, L. (2000). Bullying intervention strategies that work. *Education World*

http://www.educationworld.com/a_issues/issues103.shtml

Voors, W. (2003). Bullying – both sides of the fence, *Paradigm* Winter 2003, 8(1) & 6(4), 16.